

Contents

Chapter 1. Introduction	1
Chapter 2. Preliminaries: Tate cohomology and flabby resolutions	23
Chapter 3. CARAT ID of the \mathbb{Z} -classes in dimensions 5 and 6	29
Chapter 4. Krull-Schmidt theorem fails for dimension 5	33
4.0. Classification of indecomposable maximal finite groups $G \leq \mathrm{GL}(n, \mathbb{Z})$ of dimension $n \leq 6$	37
4.1. Krull-Schmidt theorem (1)	41
4.2. Krull-Schmidt theorem (2)	50
4.3. Maximal finite groups $G \leq \mathrm{GL}(n, \mathbb{Z})$ of dimension $n \leq 6$	59
4.4. Bravais groups and corresponding quadratic forms	61
Chapter 5. GAP algorithms: the flabby class $[M_G]^{fl}$	79
5.0. Determination whether M_G is flabby (coflabby)	79
5.1. Construction of the flabby class $[M_G]^{fl}$ of the G -lattice M_G	81
5.2. Determination whether $[M_G]^{fl}$ is invertible	84
5.3. Computation of E with $[[M_G]^{fl}]^{fl} = [E]$	86
5.4. Possibility for $[M_G]^{fl} = 0$	89
5.5. Verification of $[M_G]^{fl} = 0$: Method I	93
5.6. Verification of $[M_G]^{fl} = 0$: Method II	97
5.7. Verification of $[M_G]^{fl} = 0$: Method III	105
Chapter 6. Flabby and coflabby G -lattices	113
Chapter 7. $H^1(G, [M_G]^{fl}) = 0$ for any Bravais group G of dimension $n \leq 6$	125
Chapter 8. Norm one tori	129
Chapter 9. Tate cohomology: GAP computations	133
Chapter 10. Proof of Theorem 1.27	139
Chapter 11. Proof of Theorem 1.28	153
Chapter 12. Proof of Theorem 12.3	165
Chapter 13. Application of Theorem 12.3	173

Chapter 14. Tables for the stably rational classification of algebraic k -tori of dimension 5	179
Bibliography	211