

Variational Methods for Nonlocal Fractional Problems

GIOVANNI MOLICA BISCI

Università 'Mediterranea' di Reggio Calabria, Italy

VICENTIU D. RADULESCU

Institute of Mathematics of the Romanian Academy

RAFFAELLA SERVADEI

Università degli Studi di Urbino 'Carlo Bo', Italy



CAMBRIDGE
UNIVERSITY PRESS

Contents

<i>Foreword</i>	xiii
<i>Preface</i>	xv
Part I Fractional Sobolev spaces	
1 Fractional framework	3
1.1 Fourier transform of tempered distributions	4
1.2 Fractional Sobolev spaces	5
1.2.1 Embedding properties	7
1.2.2 The Sobolev space $H^s(\Omega)$	11
1.3 The fractional Laplacian operator	11
1.3.1 The constant $C(n, s)$: some properties	14
1.3.2 The fractional Laplacian via Fourier transform	20
1.3.3 A generalization of $(-\Delta)^s$	22
1.4 The fractional Sobolev space $\tilde{H}_0^s(\Omega)$	22
1.4.1 The extension problem and the space $X_0^s(\mathcal{C}\Omega)$	23
1.5 Other fractional Sobolev-type spaces	26
1.5.1 The space $X_0^s(\Omega)$	36
1.5.2 Embedding properties of $X_0^s(\Omega)$	40
1.5.3 The general case	41
2 A density result for fractional Sobolev spaces	43
2.1 The main theorems	44
2.2 Some preliminary lemmas	46
2.2.1 Properties of the support	46
2.2.2 Acting by convolution	48
2.2.3 Cutoff technique	50
2.2.4 Effect of the translations	52

2.3	Proof of Theorem 2.2 (and of Remark 2.3)	53
2.4	Proof of the main result	55
2.5	Note on the partition of unity	59
3	An eigenvalue problem	63
3.1	Eigenvalues and eigenfunctions of $-\mathcal{L}_K$	64
3.2	A direct approach	65
3.2.1	Proof of Proposition 3.1	68
3.3	Another variational characterization of the eigenvalues	77
3.4	A regularity result for the eigenfunctions	78
3.5	Nodal set of the eigenfunctions of $(-\Delta)^s$: 1D case	82
4	Weak and viscosity solutions	84
4.1	Viscosity solutions	84
4.2	Perron method and existence theory for viscosity solutions	86
4.3	Regularity theory for weak solutions	88
4.3.1	Maximum principle for weak solutions	90
4.3.2	Proof of Theorem 4.2	90
4.4	Proof of Theorem 4.1	93
4.5	On the boundedness of weak solutions	95
4.5.1	The linear case	95
4.5.2	The nonlinear case	100
5	Spectral fractional Laplacian problems	104
5.1	Two different fractional operators	104
5.2	A comparison between the eigenfunctions of A_s and $(-\Delta)^s$	106
5.2.1	Poisson kernel of fractional type	107
5.2.2	Optimal regularity for the eigenfunctions of $(-\Delta)^s$	108
5.3	The spectrum of A_s and $(-\Delta)^s$	117
5.4	One-dimensional analysis	117
5.5	The first eigenvalue of A_s and that of $(-\Delta)^s$	124
Part II Nonlocal subcritical problems		
6	Mountain pass and linking results	131
6.1	Hypotheses and statements	131
6.2	Estimates on the nonlinearity and its primitive	133
6.3	Proofs of the main theorems	135
6.3.1	The case $\lambda < \lambda_1$: mountain pass-type solutions	136
6.3.2	The case $\lambda \geq \lambda_1$: linking-type solutions	143
6.4	Comments on the sign of the solutions	148
6.5	A remark on the case $\lambda = 0$	150

7	Existence and localization of solutions	152
7.1	Existence of one weak solution	153
7.1.1	Notation	154
7.1.2	Main results	155
7.1.3	The fractional Laplacian setting	157
7.2	A doubly parametric problem	161
7.2.1	Some bifurcation theorems	162
7.2.2	Proof of Theorem 7.10	164
8	Resonant fractional equations	169
8.1	A saddle point result	169
8.2	Eigenvalues for linear problems with weights	171
8.3	Some technical lemmas	173
8.4	The main result	177
8.4.1	Geometry of the functional \mathcal{J}_a	178
8.4.2	The Palais–Smale condition	180
8.4.3	Proof of Theorem 8.1	184
9	A pseudoindex approach to nonlocal problems	186
9.1	A multiplicity result	186
9.1.1	Variational formulation	188
9.2	A pseudoindex theorem	188
9.3	The Palais–Smale condition	190
9.4	Some preparatory lemmas	192
9.5	$k - h + 1$ distinct pairs of solutions	194
10	Multiple solutions for parametric equations	195
10.1	Two abstract critical points results	195
10.2	Three weak solutions	197
10.3	Two weak solutions	201
11	Infinitely many solutions	206
11.1	The main results	207
11.2	Abstract approach	209
11.2.1	Some preliminary lemmas	211
11.3	Some compactness conditions	213
11.3.1	The Palais–Smale condition for $\mathcal{J}_{K,\lambda,h}$	213
11.3.2	The Palais–Smale condition on M for $\mathcal{J}_{K,\lambda,h}$	216
11.4	Existence of infinitely many solutions	220
11.4.1	Proof of Theorem 11.2	220

12	Fractional Kirchhoff-type problems	224
12.1	Nondegenerate Kirchhoff equations	225
12.1.1	Mountain pass solution	225
12.1.2	Multiple solutions	227
12.1.3	Kirchhoff equations with bounded primitive	227
12.2	Degenerate Kirchhoff equations	231
12.2.1	Some multiplicity results	231
12.2.2	A Clark-type result for a Kirchhoff model	236
13	On fractional Schrödinger equations	240
13.1	The main problem	241
13.1.1	Assumptions on the potential and the nonlinearity	241
13.1.2	The abstract framework	242
13.2	Multiple solutions	244
13.3	Nonexistence results	245
13.4	Perturbed Schrödinger equations	247
Part III Nonlocal critical problems		
14	The Brezis–Nirenberg result for the fractional Laplacian	251
14.1	A critical fractional Laplace equation	252
14.2	Geometry of the functional $\mathcal{J}_{s,\lambda}$	255
14.3	Some crucial estimates	256
14.3.1	Some remarks on condition $S_{s,\lambda} < S_s$	267
14.4	End of the proof of Theorem 14.1	270
15	Generalization of the Brezis–Nirenberg result	276
15.1	Main results	276
15.1.1	Strategy for proving Theorem 15.1	277
15.2	A local Palais–Smale condition for the functional $\mathcal{J}_{s,\lambda}$	278
15.3	The geometry of the functional $\mathcal{J}_{s,\lambda}$	282
15.4	A nondegeneracy estimate	284
15.5	Proof of Theorem 15.1	289
15.5.1	A comment on the estimate of the critical level	290
15.5.2	End of the proof of Theorem 15.1	291
16	The Brezis–Nirenberg result in low dimension	293
16.1	Existence of a nontrivial solution	293
16.2	Proof of Theorem 16.1	294
16.2.1	Estimates of the critical level of $\mathcal{J}_{s,\lambda}$	295
16.2.2	End of the proof of Theorem 16.1	298

17	The critical equation in the resonant case	299
17.1	Main results	299
17.2	Estimate of the minimax critical level	301
18	The Brezis–Nirenberg result for a general nonlocal equation	309
18.1	Assumptions and main results	309
18.2	Some preliminary results	312
18.2.1	Estimates on the nonlinearity	312
18.2.2	Variational formulation of the problem	313
18.3	The critical case with a lower-order perturbation	314
18.3.1	End of the proof of Theorem 18.1	316
18.4	The critical case $\mathcal{L}_K u + \lambda u + u ^{2_s^* - 2} u = 0$	322
18.4.1	End of the proof of Theorem 18.2	323
18.5	The general case $\lambda > 0$	323
18.5.1	Some comments on the main theorems	324
18.5.2	The Palais–Smale condition for the functional $\mathcal{J}_{K,\lambda,f}$	325
18.5.3	The geometry of the functional $\mathcal{J}_{K,\lambda,f}$	331
18.5.4	Estimates of the critical level of $\mathcal{J}_{K,\lambda,f}$	334
18.6	The model case	335
19	Existence of multiple solutions	337
19.1	A multiplicity result	338
19.2	Proof of Theorem 19.1	339
20	Nonlocal critical equations with concave-convex nonlinearities	344
20.1	Main results	345
20.1.1	Variational formulation of the problem	345
20.2	The critical and concave case $0 < q < 1$	346
20.2.1	The Palais–Smale condition for $\tilde{\mathcal{F}}_{s,\lambda,q}$	350
20.2.2	Proof of statement (d) of Theorem 20.1	359
20.3	The critical and convex case $q > 1$	363
20.3.1	The Palais–Smale condition for $\mathcal{J}_{s,\lambda,q}$	363
20.3.2	Proof of Theorem 20.2	367
	<i>Bibliography</i>	371
	<i>Index</i>	381